

AD-A052 667

TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY
THE SYNTHESIS AND STRUCTURE OF A NOVEL DIMETALLIC TETRAAZANULE--ETC(U)
MAR 78 M TSUTSUI, R L BOBSEIN, R PETTERSEN

F/G 7/3

N00014-75-C-0417

UNCLASSIFIED

TR-23

NL

| OF |
AD
A052667



END
DATE
FILMED
5-78

DDC

AD A 052667

AD No. 1
DDC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract ¹⁵ ~~NO 0014-75-C-0417~~

Task No. NR 053-559

⁹ Technical Report, No. 23

12

⁶ The Synthesis and Structure of a Novel Dimetallic Tetraazaanulene:
[Dibenzo[b,i][1,4,8,11]Tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)]
micron

by

¹⁰ M. Tsutsui, R. L. Bobsein, R. Pettersen ~~and~~ R. Haaker

Prepared for Publication

in the

Journal of Inorganic Chemistry

Texas A&M University
Department of Chemistry
College Station, Texas 77843

¹¹ 15 March ~~1975~~ 1978

¹² 13P.

¹⁴ TR-23

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

47 DDC
RECEIVED
MAR 30 1978
D

407 997

JOB

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 23	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Synthesis and Structure of a Novel Dimetallic Tetraazaannulene: μ -[Dibenzo[b,i][1,4,8,11]Tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)]	5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s) M. Tsutsui, R. L. Bobsein, R. Pettersen and R. Haaker	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0417	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Texas A&M University College Station, Texas 77843	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-559	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217	12. REPORT DATE March 15, 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 10	
	15. SECURITY CLASS. (of this report)	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in the Journal of Inorganic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Dimetallic Tetraazaannulene μ -[Dibenzo[b,i][1,4,8,11]Tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)] Dibenzotetraannulene		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The first dinuclear metallotetraazaannulene complex, μ -[dibenzo[b,i][1,4,8,11]tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)], with an unusual geometry was synthesized by reaction of dirhenium decarbonyl and 1,8-dihydrodibenzo[b,i][1,4,8,11]tetraazaannulene. This compound was characterized via visible, infrared, and mass spectral methods as well as elemental analysis. Structural data were obtained by single crystal x-ray diffraction. The distance between the rhenium atoms is 3.345(1) Å. This is a significant increase (.219(1) Å) from the corresponding porphyrin compounds, the dibenzotetraazaannulene ring is puckered a significant amount from planarity.		

DD FORM 1473

1 JAN 73

EDITION OF 1965 IS OBSOLETE

S/N 0102- LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

The Synthesis and Structure of a Novel Dimetallic Tetraazaannulene:
 μ -[Dibenzo[b,i][1,4,8,11]Tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)]

by

M. Tsutsui, R. L. Bobsein, R. Pettersen and R. Haaker
Texas A&M University
Department of Chemistry
College Station, Texas 77843

ABSTRACT

The first dinuclear metallotetraazaannulene complex, μ -[dibenzo-[b,i][1,4,8,11]tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)], with an unusual geometry was synthesized by reaction of dirhenium decacarbonyl and 1,8-dihydrodibenzo[b,i][1,4,8,11]tetraazaannulene. This compound was characterized via visible, infrared, and mass spectral methods as well as elemental analysis. Structural data were obtained by single crystal x-ray diffraction. The distance between the rhenium atoms is 3.345(1) Å. This is a significant increase (.219(1) Å) from the corresponding porphyrin compounds, the dibenzotetraazaannulene ring is puckered a significant amount from planarity.


ACCESSION for	
WTLG	White Section <input checked="" type="checkbox"/>
DDC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. AND/OR SPECIAL
A	

DDC
RECEIVED
MAR 30 1978
D

The Synthesis and Structure of a Novel Dimetallic Tetraazaannulene:
 μ -[Dibenzo[b,1][1,4,8,11]tetraaza[14]annulenato]-bis[tricarbonyl rhenium(I)]

Sir:

We would like to report the synthesis and x-ray diffraction analysis of the first dinuclear metallotetraazaannulene in which one tetraazaannulene ligand binds with 2 metal atoms. There is growing interest in synthetic macrocyclic ligands because of their obvious similarity with porphyrins which have importance as biological models¹, because of their catalytic ability², and because of their unique nature in coordinating metals.³⁻¹² Homodinuclear complexes of porphyrins have been well studied.¹³⁻²² To our knowledge, no stable homodinuclear metallo-tetraazaannulenes with this novel geometry have been prepared. Such a compound would prove useful in comparing the effect of the central 'hole' or 'core' size of the macrocyclic ligand on the rhenium-rhenium distance as well as on the degree of puckering of the ligand. We have synthesized μ -[dibenzo[b,1][1,4,8,11]tetraaza[14]annulenato]-bis-[tricarbonyl rhenium(I)] (I), and we have obtained crystals suitable for x-ray study.



The compound I was prepared by refluxing dirhenium decacarbonyl²³ in equimolar quantities with 1,8-dihydrodibenzo[b,1][1,4,8,11] tetraaza-[14]annulene²⁴ in refluxing decalin for 24 hours under argon. After cooling, the resulting precipitate was centrifuged out and washed with decalin and hexane. The washed precipitate was then introduced into a neutral alumina column using chloroform as the elutant. Compound I, the wine-red fraction, was collected and evaporated to dryness (36% yield). The red residue was crystallized from a dichloromethane-chloroform-cyclohexane solution to give ruby-red crystals of I, m.p. 278-279°C. Analysis²⁵ calculated for $\text{Re}_2\text{C}_{24}\text{H}_{14}\text{N}_4\text{O}_6$: C, 34.87; H, 1.70; N, 6.78; Re, 45.04. Found: C, 34.62; H, 2.27; N, 6.64; Re, 46.03. The visible spectrum of I in chloroform has a broad absorption band centered at 455nm. ($\log \epsilon$ 3.52). The infrared spectrum of I in the solid phase (KBr) shows two strong metal-carbonyl bands at 2020 and 1900 cm^{-1} as well as the 739 $^{-1}$ band characteristic of the tetraazaannulene ligand. The mass spectrum²⁶ of I shows the parent peak at m/e 829.9 (M^+). In addition, the $(M-2\text{CO})^+$, $(M-3\text{CO})^+$, $(M-4\text{CO})^+$, $(M-5\text{CO})^+$, and $(M-6\text{CO})^+$ peaks are also observed. Since naturally occurring rhenium has two common isotopes, 185 Re and 187 Re, the triplet at $(M)^+$, $(M+2)^+$, and $(M+4)^+$ as well as this same type of triplet pattern at the major peaks gives further evidence that there are two rhenium atoms per molecule.

Crystals of I are triclinic, space group $B\bar{1}$ with $a = 10.012(3)$, $b = 4.550(1)$, $c = 13.259(4)$ Å, $\alpha = 72.92(2)$, $\beta = 93.91(2)$, $\gamma = 92.67(2)$ and $V = 575.8(3)$ Å³; $Z = 1$ for $D_{\text{meas}} = 2.36 \text{ g}\cdot\text{cm}^{-3}$ and $D_{\text{calc}} = 2.38 \text{ g}\cdot\text{cm}^{-3}$.

The data crystal dimensions are 0.11x0.14x0.09 mm, with $\mu = 111.6 \text{ cm}^{-1}$ (Mo K α : $\lambda = 0.71069 \text{ \AA}$). A total of 1382 independent reflection intensities was measured by the stationary crystal/stationary counter method using a manual General Electric XRD-700 diffractometer equipped with balanced Zr/Y filters ($2\theta_{\text{max}} = 60.00^\circ$). Ten second counts were recorded for each filter. The intensity data were processed by calculations described previously.²⁷ An experimentally determined absorption correction was applied to the data as a function of the diffractometer setting angles of ϕ and 2θ . The points were measured by rotating the crystal about the ϕ axis at the respective reflection 2θ angles with χ set at 90.00° . A maximum of 20% was noted for intensity variation. 1083 of the reflections were classed to be significantly above background.

The correct phase angles were determined by the heavy-atom technique. Atom coordinates and anisotropic temperature factors were refined to convergence by full-matrix least-squares calculations to $R = 0.020$ and $R_w = 0.022$. Hydrogen atoms could not be located in a difference Fourier map, but their scattering contribution was included in the final cycles of refinement by assuming idealized C-H geometry. Hydrogen atom parameters were not allowed to refine. The largest residual peak in a final difference map is $0.50 \text{ e} \cdot \text{\AA}^{-3}$, located 0.7 \AA from the rhenium atom. Most of the crystallographic calculations were done with the SHELEX system of programs.²⁸

The molecule lies on a crystallographic center of symmetry, and provides confirmation of the centrosymmetric space group. The figure is an ORTEP stereographic view of the molecular structure. The Table lists some interatomic distances with estimated standard deviations.

Some pertinent torsional angles are listed to show the non-planarity of the dibenzotetraazaannulene ligand.

Table*

Interatomic Distances

Re-N1	2.130(6) Å	N1-C1	1.30(1) Å
Re-N2	2.232(6)	N1-C4'	1.45(1)
Re-N1'	3.325(7)	N2-C3	1.42(1)
Re-N2'	2.334(6)	N2-C9	1.45(1)
Re-Re'	3.345(1)	C1-C2	1.48(1)
Re-C10	1.910(8)	C2-C3	1.34(1)
Re-C11	1.910(8)	C4-C5	1.38(1)
Re-C12	1.916(8)	C4-C9	1.39(1)
C10-O10	1.16(1)	C5-C6	1.42(1)
C11-O11	1.15(1)	C6-C7	1.38(1)
C12-O12	1.14(1)	C7-C8	1.37(1)
N1-N2	2.327(8)	C8-C9	1.39(1)
N1-N2'	2.615(8)		

Torsional Angles

N1'-C4-C9-N2	-1.4(1.1)°	C3-C2-C1-N1	27.2(1.2)°
C4-C9-N2-C3	-153.8(8)	C2-C1-N1-C4'	-157.2(6)
C9-N2-C3-C2	-170.2(8)	C1-N1-C4'-C9'	131.7(8)
N2-C3-C2-C1	-10.8(1.3)		

* The prime (') after an atom label indicates the atom generated by the inversion symmetry operation.




Figure 1. An ORTEP stereographic view of the molecular structure of μ -[dibenzo[b,i][1,4,8,11]tetraaza[14]annuleno]-bis(tricarbonyl rhenium (I)) (I).

The tricarbonylrhenium(I) groups are located on opposite sides of the dibenzotetraazaannulene ligand. Interatomic distances indicate each rhenium is bonded to three nitrogen atoms and that they both bond in common to N2 and N2'. The bond distances within the organic ligand clearly show the non-conjugation of the macrocycle. The distances of nitrogen atoms N1 and N2 from the center of the ligand are 2.236(8) and 1.555(8) Å respectively. This compares with 2.26(1) and 1.81(2) Å for a similar dirhenium structure with a tetraphenylporphyrin ligand.¹³ The deviation of the rhenium atom from the best least-squares plane containing the N1, N2, N1' and N2' atoms is 1.50 Å.

The crystal lattice appears to be disordered. An ordered lattice of space group $B\bar{1}$ requires two molecules per unit cell, whereas packing considerations and a density measurement indicate one molecule per cell. The least-squares calculations were done using half-occupancy for all atoms. Systematically absent reflections ($h+1 = 2n + 1$) were carefully checked as well as possible reflections with the k index divided by two or three. No additional reflections could be found. A suitable packing model was determined by systematically removing half of the molecules in the overall lattice. Transformation to a C-face centered lattice shows that this packing model corresponds to the transformed cell with the C-face center removed. A check was made for the $h+k = 2n + 1$ reflections of the transformed cell, but these were also found to be systematically absent. No satisfactory explanation for the apparent lattice disorder has been found.

It is significant to note that the smaller 'hole' size of the dibenzotetraazaannulene ligand has caused the rhenium atoms to be farther apart ($.219(1) \text{ \AA}$) than in the porphyrin analogs. Like the porphyrin ligand in these complexes, the dibenzotetraazaannulene ligand also shows a significant amount of twisting in its complex with two rhenium tricarbonyl moieties.

During the workup of I, a second dark brown band was eluted from the neutral alumina column. It was formulated as [monohydrogen dibenzo[b,i][1,4,8,11]tetraaza[14]annulenato]tricarbonyl rhenium(I) in analogy to the porphyrin systems already known, but attempts to obtain analytical purity of this compound have failed.

We are continuing our studies of molecules of these types.

Acknowledgement. The authors wish to acknowledge support of this research project by the Office of Naval Research and The Robert A. Welch Foundation (Grant No. A-614).

REFERENCES

1. M. Tsutsui, C. P. Hsung, D. Ostfeld, T. S. Srivastava, D. L. Cullen, and E. F. Meyer, Jr., J. Amer. Chem. Soc., **97**, 3952 (1975).
2. F. Beck, W. Dammert, J. Heiss, H. Hiller, and R. Polster, Z. Naturforsch., **28**, 1009 (1973).
3. L. G. Bell and J. C. Dabrowiak, J. Chem. Soc., Chem. Commun., 512 (1975).
4. D. R. Neves and J. C. Dabrowiak, Inorg. Chem., **15**, 129 (1976).
5. V. L. Goedken, J. Molin-Case, and Y.-A. Whang, J. Chem. Soc., Chem. Commun., 337 (1973).
6. V. L. Goedken, S.-M. Peng, and Y.-A. Park, J. Amer. Chem. Soc., **96**, 204 (1974).
7. V. L. Goedken, Y.-A. Park, S.-M. Peng, and J. Norris, J. Amer. Chem. Soc., **96**, 7693 (1974).
8. V. L. Goedken, J. J. Pluth, S.-M. Peng, and B. Bursten, J. Amer. Chem. Soc., **98**, 8014 (1976).
9. V. L. Goedken and Y.-A. Park, J. Chem. Soc., Chem. Commun., 214 (1975).
10. M. C. Weiss, B. Bursten, S.-M. Peng, and V. L. Goedken, J. Amer. Chem. Soc., **98**, 8021 (1976).
11. S.-M. Peng, J. A. Ibers, M. Millar, and R. H. Holm, J. Amer. Chem. Soc., **98**, 8037 (1976).
12. S.-M. Peng and V. L. Goedken, J. Amer. Chem. Soc., **98**, 8500 (1976).
13. S. Kato, M. Tsutsui, D. L. Cullen, and E. F. Meyer, Jr., J. Amer. Chem. Soc., **99**, 620 (1977).
14. Z. Yoshida, H. Ogoshi, T. Omura, E. Watanabe, and T. Kurosaki, Tetrahedron Lett., 1077 (1972).
15. A. Takenaka, Y. Sasada, W. Ogoshi, T. Omura, and Z. Yoshida, Acta Crystallogr., Sect. B, **31**, 1 (1975).
16. M. F. Hudson and K. M. Smith, J. Chem. Soc., Chem. Commun., 515 (1973).
17. M. F. Hudson and K. M. Smith, Tetrahedron Lett., 2223 (1974).
18. P. Clare and F. Glockling, Inorganica Chimica Acta, **14** (1975).

19. D. Cullen, E. Meyer, T. S. Srivastava, and M. Tsutsui, J. Amer. Chem. Soc., 94, 7603 (1972).
20. D. Ostfeld, M. Tsutsui, C. P. Hsung, and O. C. Conway, J. Amer. Chem. Soc., 93, 2548 (1971).
21. M. Tsutsui and C. P. Hsung, J. Amer. Chem. Soc., 95, 5777 (1973).
22. M. Tsutsui and C. P. Hsung, Chem. Lett., 941 (1973).
23. $\text{Re}_2(\text{CO})_{10}$ obtained from Strem Chemicals Inc.
24. H. Hiller, P. Dimroth, and H. Pfitzner, Justus Liebigs Ann. Chem., 717, 137 (1968).
25. Elemental analyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, N. Y. 11377.
26. Center for Trace Characterization, Texas A&M University, College Station, Tx. 77843.
27. R. C. Pettersen and H. L. Kim, J. Chem. Soc. , Perk.II, 1399 (1975).
28. SHELX, 1976: A program system written by G.M. Sheldrick, University of Cambridge, U.K.




Figure 1.

M. Tsutsui and others

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market St., Rm. 447 San Francisco, Calif. 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Dr. R. M. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1
Commander Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1
		Dr. C. Quicksall Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1

TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1		
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1		
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1		
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1		
Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1		
Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1		
Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1		